

Treating Your Non-Small Cell Lung Cancer (NSCLC): Start a Conversation With Your Doctor

How this guide can help

Living with NSCLC—and the changes that come with it—can be overwhelming. These feelings are normal, and this guide is here to help. Within, you'll find helpful thought-starters and questions to discuss with your doctor or healthcare team. Using these prompts can help you make the most out of your discussions so you can make informed decisions throughout your treatment journey.

Here are a few helpful tips to make the most of your appointments:



Write down any questions you have ahead of time



Circle questions within this guide that you'd like to discuss



Bring a notebook or use the space in this guide to take notes during your conversation



Bring a family member or friend with you to help take notes during your conversation

What is EXKIVITY?

EXKIVITY is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC):

- that has spread to other parts of the body (metastatic) and cannot be removed by surgery, **and**
- has a certain abnormal epidermal growth factor receptor (*EGFR*) gene, **and**
- whose disease has worsened while on or after chemotherapy that contains platinum

Your healthcare provider will perform a test to make sure that EXKIVITY is right for you. It is not known if EXKIVITY is safe and effective in children.

EXKIVITY is approved based on a medical study that measured how many people responded to treatment and for how long. There are ongoing studies to confirm its continued approval.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about EXKIVITY?

EXKIVITY may cause serious side effects, including changes in the electrical activity of your heart called QTc prolongation and Torsades de Pointes. QTc prolongation can cause irregular heartbeats that can be life-threatening and may lead to death. Your healthcare provider will check the electrical activity of your heart with a test called an electrocardiogram (ECG) and do blood tests to check your electrolytes before starting treatment and during treatment with EXKIVITY. Tell your healthcare provider right away if you feel dizzy, lightheaded, faint or have an irregular heartbeat.

Please read the Important Safety Information including information for the Boxed Warning on pages 6-7 and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

 **EXKIVITY**[®]
mobocertinib
40 mg capsules

Preparing for Your Conversation

The better prepared you are to discuss your diagnosis and treatment options with your doctor, the more productive the conversation will be. Consider asking some of the questions below to assess where you are in your treatment journey and to find out what may come next.



Your diagnosis

- What is the stage of my lung cancer?
- What does *metastatic* mean?
- What does it mean for cancer to progress?
- What biomarker testing was done to determine my type of lung cancer?
- Can you explain the results of my biomarker testing?
- What is an epidermal growth factor receptor (*EGFR*) Exon20 insertion mutation? How does it apply to my cancer?



Your treatment goals

- What is the overall goal of my treatment?
- What is most important when deciding on a treatment?

Check what's most important to you:

How long people respond to treatment

Manageable side effects

Likelihood of treatment working for me

Cost of treatment

Ability to take treatment at home

Other _____

- What are all the treatment options for my type of cancer?
- What treatment plan do you recommend? Why?
- How will we know if treatment is working?
- Would a clinical study be right for me?
- Should I get a second opinion? If so, can you recommend someone?
- What's next if my treatment doesn't work?



Getting support

- How do we reach you over holidays, on weekends, or after office hours?
- What support services are available to me and my family?
- Can you recommend a social worker to help me find support services?
- What advocacy groups are available to help support me throughout my journey?
- Are there any financial assistance programs to help with the cost of my treatment?

Please read the Important Safety Information including information for the Boxed Warning on pages 6-7 and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

 **EXKIVITY**[®]
mobocertinib
40 mg capsules

Having a Conversation About EXKIVITY® (mobocertinib)

One treatment option you may want to talk to your doctor about is EXKIVITY.

EXKIVITY is the first oral therapy designed for patients with epidermal growth factor receptor (*EGFR*) Exon20 insertion-positive metastatic non-small cell lung cancer (NSCLC) after chemotherapy that contains platinum. Use the questions below to learn more about EXKIVITY, and to see if it's right for you.



Taking EXKIVITY

- How is EXKIVITY taken?
- How often do I need to take it?
- Do I need to take EXKIVITY with food?
- For how long will I need to take EXKIVITY?
- Will I need to change what I eat or drink during treatment?
- Will I need any routine tests or scans before taking EXKIVITY? What about while I'm on EXKIVITY? How often will they take place?
- How soon can I start treatment?



Clinical study results with EXKIVITY

- What results have other people seen with EXKIVITY?
- How long are people normally treated with EXKIVITY?
- How will we know if EXKIVITY is working for me?
- How soon will I see results with EXKIVITY?



Side effects with EXKIVITY

- What are the common side effects of EXKIVITY, and what should I look out for?
- Is there anything I can do to prepare for common side effects?
- What can we do to help manage side effects if they occur?
- Whom should I reach out to if I experience any side effects?
- What should I do if I experience one of the common side effects, such as diarrhea or skin reactions?



Support for people taking EXKIVITY

- How can I find out if EXKIVITY is covered by my insurance? What if I don't have insurance?
- Does EXKIVITY have a patient support program?

Please read the Important Safety Information including information for the Boxed Warning on pages 6-7 and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

 **EXKIVITY**[®]
mobocertinib
40 mg capsules

Glossary

Here's a quick reference for some words you may hear while talking with your doctor.

Immunotherapy: A type of treatment that helps your immune system fight cancer.

Infusion: A way to administer medicine into your body using a thin tube called a catheter that is put into a vein in your hand or forearm using a needle. The catheter is connected by a tube to a plastic bag that contains the medicine. A typical infusion can last anywhere from a few minutes to a few hours.

Median: The middle number in a list sorted from lowest to highest. Half the numbers in the list are the same as the median or lower, and half are the same or higher.

Oral therapy: A type of treatment that is given by mouth in tablet, capsule, or liquid form.

Targeted therapy: A type of treatment that can slow the growth, progression, and spread of cancer by targeting cancer cells or specific molecules that are responsible for cancer cell growth.

Tyrosine kinase inhibitor (TKI): A type of treatment that blocks an enzyme in your cells called a tyrosine kinase. Some cancer cells have too many of these enzymes or have enzymes that are too active. Blocking them with a TKI may help keep the cancer cells from growing.

Please read the Important Safety Information including information for the Boxed Warning on pages 6-7 and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

 **EXKIVITY**[®]
mobocertinib
40 mg capsules

Important Safety Information for EXKIVITY[®] (mobocertinib)

What is the most important information I should know about EXKIVITY?

EXKIVITY may cause serious side effects, including changes in the electrical activity of your heart called QTc prolongation and Torsades de Pointes. QTc prolongation can cause irregular heartbeats that can be life-threatening and may lead to death. Your healthcare provider will check the electrical activity of your heart with a test called an electrocardiogram (ECG) and do blood tests to check your electrolytes before starting treatment and during treatment with EXKIVITY. Tell your healthcare provider right away if you feel dizzy, lightheaded, faint or have an irregular heartbeat.

EXKIVITY may cause serious side effects, including:

- **Lung problems.** EXKIVITY may cause severe lung problems that may lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your healthcare provider right away if you develop any new or worsening symptoms, including trouble breathing or shortness of breath, cough, chest pain, or fever.
- **Heart problems, including heart failure.** EXKIVITY may cause heart problems that may lead to death. Your healthcare provider should check your heart function before you start and during treatment with EXKIVITY. Tell your healthcare provider right away if you have any signs or symptoms of a heart problem, including feeling like your heart is pounding or racing, shortness of breath, chest pain, swelling of your ankles and feet, or feeling faint.
- **Diarrhea.** Diarrhea is common during treatment with EXKIVITY and may sometimes be severe. Diarrhea can cause you to lose too much body fluid (dehydration) and kidney problems. Your healthcare provider may tell you to start drinking more fluids and electrolytes to replace body salts or start taking your antidiarrheal medicines. Tell your healthcare provider right away if you have any loose stools or have stools more often than is normal for you.

The most common side effects of EXKIVITY include diarrhea, rash, nausea, mouth sores, vomiting, decreased appetite, infection of skin around nails, tiredness, dry skin, muscle or bone pain. EXKIVITY may affect fertility in females and males, which may affect your ability to have a child. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of EXKIVITY. Call your doctor for medical advice about side effects. You may report side effects to FDA at **1-800-FDA-1088**.

Before taking EXKIVITY, tell your healthcare provider about all your medical conditions, including if you:

- have heart problems, including a condition called long QTc syndrome
- have problems with your electrolytes, such as sodium, potassium, calcium or magnesium
- have lung or breathing problems other than lung cancer
- are pregnant or plan to become pregnant. EXKIVITY can harm your unborn baby

Please read the Important Safety Information including information for the Boxed Warning on pages 6-7 and the [Patient Information](#) in the accompanying full [Prescribing Information](#).

 **EXKIVITY[®]**
mobocertinib
40 mg capsules

Important Safety Information for EXKIVITY (Cont'd)

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with EXKIVITY. You should use an effective form of non-hormonal birth control during treatment and for **1 month** after your last dose of EXKIVITY. Birth control pills (oral contraceptives) and other hormonal forms of birth control may not work as well during treatment with EXKIVITY. Talk to your healthcare provider about birth control methods that might be right for you during this time. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with EXKIVITY.

Males who have female partners who are able to become pregnant:

- You should use effective birth control during treatment and for **1 week** after your last dose of EXKIVITY.
- are breastfeeding or plan to breastfeed. It is not known if EXKIVITY passes into your breast milk. Do not breastfeed during treatment and for **1 week** after your last dose of EXKIVITY.

Tell your healthcare provider about all the medicines you take, including prescription medicines and over-the-counter medicines, vitamins, and herbal supplements. Tell your healthcare provider if you take medicines for heart problems. EXKIVITY and other medicines may affect each other causing serious side effects.

How should I take EXKIVITY?

- Take EXKIVITY exactly as your healthcare provider tells you to take it.
- Take your prescribed dose of EXKIVITY 1 time each day.
- Take EXKIVITY with or without food.
- Swallow EXKIVITY capsules whole. Do not open, chew, or dissolve the contents of the capsules.
- **Do not** change your dose or stop taking EXKIVITY unless your healthcare provider tells you to.
- Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with EXKIVITY if you develop certain side effects.
- If you miss a dose of EXKIVITY, and it has been more than 6 hours, skip the dose and take your next dose at your regularly scheduled time the next day.
- If you vomit a dose of EXKIVITY, do not take an extra dose. Take your next dose at your regularly scheduled time the next day.

What should I avoid while taking EXKIVITY?

- Avoid eating grapefruit or drinking grapefruit juice during treatment with EXKIVITY. Grapefruit may increase the amount of EXKIVITY in your blood.

Please see [Patient Information](#) in the accompanying EXKIVITY (mobocertinib) full [Prescribing Information](#).

Please read the Important Safety Information including information for the **Boxed Warning** on pages 6-7 and the [Patient Information](#) in the accompanying full [Prescribing Information](#).



ONCOLOGY

All trademarks are property of their respective owners.
©2022 Takeda Pharmaceuticals U.S.A., Inc.
All rights reserved. 04/22 USO-EXK-0052

 **EXKIVITY**[®]
mobocertinib
40 mg capsules